

Effect of Er,Cr:YSGG Laser Root Conditioning on the Success of Root Coverage with Subepithelial Connective Tissue Graft: A Randomized Clinical Trial with a 6-Month Follow-Up

Banafsheh Poormoradi ¹, Parviz Torkzaban ², Leila Gholami ¹, Amirarsalan Hooshyarfard ³✉, Maryam Farhadian ⁴

¹ Assistant Professor, Department of Periodontics, School of Dentistry, Hamadan University of Medical Sciences, Hamadan, Iran

² Associate Professor, Department of Periodontics, School of Dentistry, Hamadan University of Medical Sciences, Hamadan, Iran

³ Postgraduate Student, Department of Periodontics, School of Dentistry, Hamadan University of Medical Sciences, Hamadan, Iran

⁴ Assistant Professor, Department of Biostatistics, School of Public Health and Research Center for Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

Abstract

Objectives: Finding predictable approaches for root surface biomodification is an important challenge in the treatment of gingival recession. This study sought to assess the root coverage percentage by subepithelial connective tissue graft (SCTG) following root surface conditioning with erbium, chromium: yttrium scandium gallium garnet (Er,Cr:YSGG) laser.

Materials and Methods: In this split-mouth, randomized clinical trial, 30 teeth with Miller's Class I and II gingival recession were treated with SCTG (the Langer and Langer technique) with (case group) or without (control group) root surface conditioning with Er,Cr:YSGG laser (wavelength=2780 nm, power=0.75 W, H mode, repetition rate=20 Hz). Recession depth (RD), recession width (RW), clinical attachment level (CAL), and probing depth (PD) were assessed at the baseline (one week before surgery) and at 2 and 6 months postoperatively. The amount of root coverage was quantified in the two groups. Data were analyzed using Friedman test and Wilcoxon signed-rank test.

Results: No significant difference was noted between the case and control groups in any parameter ($P>0.05$). Significant improvement occurred in all the measured parameters in the two groups after surgery ($P<0.05$). The mean root coverage at the end of the study period was 87% and 80% in the case and control groups, respectively ($P=0.244$), and complete root coverage was achieved in 66% and 60% of the samples in the case and control groups, respectively.

Conclusions: Root surface conditioning by Er,Cr:YSGG laser improved the mean root coverage and the percentage of complete root coverage. However, these changes were not statistically significant.

Keywords: Gingival Recession; Tooth Root; Tissue Transplantation; Lasers

Journal of Dentistry, Tehran University of Medical Sciences, Tehran, Iran (2018; Vol. 15, No. 4)

✉Corresponding author:

A. Hooshyarfard, Department of Periodontics, School of Dentistry, Hamadan University of Medical Sciences, Hamadan, Iran

Dr.A.Hooshyarfard@gmail.com

Received: 15 January 2018

Accepted: 11 June 2018

INTRODUCTION

Gingival recession is an unfavorable clinical condition characterized by the migration of the gingival margin from the cemento-enamel junction (CEJ) towards the apex [1]. It results in denuding of the root surface and subsequent tooth hypersensitivity, root caries, and esthetic problems [2-4].

A high percentage of adults suffer from gingival

recession [5]. Inflammatory periodontal disease, shortage of keratinized tissue, mechanical trauma, orthodontic movement, buccal positioning of the root, bone dehiscence, and abnormal frenal attachment are among the causes of gingival recession [5].

The main goal of treatment is to cover the denuded root surface to decrease tooth hypersensitivity and improve esthetics. Different

therapeutic approaches suggested for this condition include surgical and non-surgical treatment modalities. Surgical approaches include free gingival graft [6], subepithelial connective tissue graft (SCTG) [7], coronally advanced flap [8], laterally sliding flap [9], double papillae flap [10], guided tissue regeneration [11], and acellular dermal matrix allograft [12]. The success rate of these procedures depends on several factors such as the position of the tooth, the class of recession, surgeon's experience and expertise, the surgical technique, and postoperative care [6-14]. Of the aforementioned procedures, SCTG has shown a high success rate and optimal predictability [13-15]. Thus, SCTG is considered the gold standard for the assessment of novel approaches [16].

Several studies have evaluated the efficacy of root surface conditioning aiming to improve the treatment results. It has been reported that mechanical debridement preserves the smear layer on the root surface and thus, prevents cell reattachment to this surface, compromising the process of regeneration and repair [17]. Several strategies have been proposed to overcome this problem including the use of root surface conditioners such as citric acid [18], ethylenediaminetetraacetic acid (EDTA) [19], tetracycline hydrochloride [20], and hydrogen peroxide [21], enamel matrix proteins [22], platelet-rich plasma [23], and recombinant human growth factors [24].

A recent study showed that laser irradiation eliminates the smear layer and exerts bactericidal effects and can therefore improve the condition of root surface for connective tissue attachment [25]. In contrast to carbon dioxide (CO₂) and neodymium-doped yttrium aluminum garnet (Nd:YAG) lasers that have limited applications only for the soft tissue, erbium-doped yttrium aluminium garnet laser (Er:YAG; 2940 nm) and erbium, chromium: yttrium scandium gallium garnet (Er,Cr:YSGG; 2780 nm) laser are suitable for use on hard tissues [26]. To date, published

information about the clinical outcome of the application of Er,Cr:YSGG laser for the treatment of gingival recession is scarce. Thus, this study aimed to assess and compare the clinical results of gingival recession treatment by SCTG with/without root surface conditioning with Er,Cr:YSGG laser.

MATERIALS AND METHODS

This split-mouth, randomized clinical trial has been approved by the ethics committee of Hamadan University of Medical Sciences (IR.UMSHA.REC.1396.115) and is registered at the Iranian Registry of Clinical Trials (IRCT201705309014N167).

Patient selection:

The patients were selected from among those presenting to the Periodontics Department of School of Dentistry, Hamadan University of Medical Sciences with gingival recession defects. The study was thoroughly explained to the patients, and written informed consent was obtained from them.

The inclusion criteria:

1. Good oral hygiene (plaque index <30%)
2. Miller's Class I and II gingival recession defects [27]
3. No tooth mobility
4. Absence of trauma from occlusion

The exclusion criteria:

1. Positive medical history contraindicating dental interventions
2. Presence of coagulation problems
3. Intake of medications interfering with periodontal health or the healing process
4. Alcohol consumption, tobacco use, or cigarette smoking
5. Disability or not showing up for follow-up sessions

Scaling, root planing, and crown polishing were performed for all patients four weeks prior to surgery. Oral hygiene instructions were also given to the patients. To ensure the absence of periapical lesions, a parallel periapical

radiograph was taken from the respective teeth, and the teeth were then randomly divided into two groups of case and control.

Assessment of clinical parameters:

The following clinical parameters were measured at the buccal surface of the teeth one week prior to surgery and at 2 and 6 months postoperatively using a periodontal probe (Williams Periodontal Probe, Hu-Friedy, Chicago, IL, USA):

- Recession depth (RD): From the CEJ to the lowest point of migration of the gingival margin.
- Recession width (RW): Distance from the mesial to the distal aspect of the gingival margin at the level of the CEJ.
- Clinical attachment level (CAL): From the CEJ to the bottom of the gingival sulcus.
- Probing depth (PD): From the gingival margin to the bottom of the gingival sulcus.

Calibration of the examiner:

The intraclass correlation coefficient was calculated for the assessment of the reproducibility of measurements, which was found to be 0.99, and indicated an excellent intraobserver agreement.

Treatment protocol:

For all patients, 10% povidone-iodine (Iran Najo Pharmaceutical Hygienic & Cosmetic Co., Tehran, Iran) and 0.2% chlorhexidine (Iran Najo Pharmaceutical Hygienic & Cosmetic Co., Tehran, Iran) were used for extraoral and intraoral disinfection, respectively. Lidocaine plus epinephrine (Persocaine-E®, Darou Pakhsh Pharmaceutical Mfg. Co., Tehran, Iran) was used for local anesthesia.

Gingival recession defects were treated according to the Langer and Langer technique [7]. A partial-thickness flap was elevated with two vertical incisions wider than the recession area by the length or half-length of a tooth mesiodistally. The coronal margin of the flap was prepared by a sulcular horizontal incision.

Interdental papilla remained untouched. The flap was extended to the mucobuccal fold without causing any perforation. Root surfaces were curetted to eliminate irregularities and dental plaque. A proper size connective tissue graft with 2 mm thickness was harvested from the palate using the trap-door technique [14]. The area was sutured with non-resorbable stitches (Braided Silk 4-0, SUPASIL, SUPA Medical Devices Co., Tehran, Iran).

Er,Cr:YSGG laser (Waterlase; Biolase Technologies, San Clemente, CA, USA) was used for root surface conditioning in the case group. The laser optic fiber was positioned perpendicular to the surface at a distance of 1-2 mm. The laser was irradiated at 2780 nm wavelength, 20 Hz repetition rate, 0.75 W power, H mode [28] with 60% water and 40% air, using a Gold handpiece with MZ6 tip (600 µm in diameter and 6 mm in length) in spiral motion (vertical, horizontal, and oblique directions) and defocused mode. Eyeglasses with a suitable optical density were worn.

The graft was trimmed if required and was then fixed at the recipient site using resorbable stitches (Polyglycolate coated 4-0, SUPABON, SUPA Medical Devices Co., Tehran, Iran). For better blood supply, the flap covered a large part of the graft. Eugenol-free periodontal dressing (Coe-Pak, GC America, Alsip, IL, USA) was applied on the surgical site and was repeated after one week. At the end of the second week, the stitches were removed.

The root coverage treatments in each patient were performed with an interval of 6 weeks between the first and the second surgeries.

Postoperative care:

Amoxicillin (500 mg every 8 hours for 7 days; LOGHMAN Pharmaceutical & Hygienic Co., Tehran, Iran) and ibuprofen (400 mg every 6 hours for 48 hours; ADVIFEN®, ZAHRAVI Pharmaceutical Co., Tehran, Iran) were prescribed postoperatively. The patients were requested to use soft food and not to brush the

teeth at the surgical site for 14 days. Also, 0.2% chlorhexidine mouthwash was prescribed twice a day, each time for one minute. The sutures were then removed, chlorhexidine was prescribed for two more weeks, and tooth brushing with a soft toothbrush was recommended twice a day. Dental prophylaxis was performed two weeks after suture removal and then monthly until the end of the study period. To ensure the absence of bias, all surgical treatments were done by a single surgeon, whereas the clinical measurements were made by another examiner.

Figure 1 shows the stages of the treatment in a patient.

Assessment of root coverage:

The percentage of root coverage was calculated using the following formula:

$$\text{Root Coverage (RC)} = (\text{Preoperative RD} - \text{Postoperative RD}) / \text{Preoperative RD} \times 100$$

Data analysis:

Data were analyzed by SPSS version 24 software program (SPSS Inc., Chicago, IL, USA). Descriptive data were reported. Since data were not normally distributed according to Kolmogorov-Smirnov test, non-parametric tests were used for data analysis. Intragroup comparisons of the variables were made using Friedman test.



Fig. 1: (A) Tooth #12 before treatment. (B) Elevation of a partial-thickness flap. (C) Root surface conditioning. (D) Graft placement at the recipient site. (E) Suturing the flap at the recipient site. (F) Tooth #12 after treatment

In case of significant results, Wilcoxon signed-rank test with Bonferroni adjustment was applied for pairwise comparisons. Wilcoxon signed-rank test was used for intergroup comparisons. Alpha=0.05 was considered statistically significant for all comparisons, except for those requiring Bonferroni adjustment, for which, alpha=0.008 was considered significant.

RESULTS

Five male patients with a mean age of 36.4 ± 8.35 years were enrolled. Thirty teeth requiring treatment for gingival recession were evaluated in two groups of 15, matched in terms of the type of the tooth (anterior teeth or premolars) and the type of recession (Miller's Class I or II). Both case and control groups included 5 anterior teeth with Miller's Class I recession, 2 anterior teeth with Miller's Class II recession, 3 premolars with Miller's Class I recession, and 5 premolars with Miller's Class II recession.

The clinical parameters were compared at the baseline (one week before surgery) and at 2 and 6 months postoperatively between the two groups (Table 1).

No significant difference was found between the two groups at different time points ($P > 0.05$). In the case group, significant changes occurred in RD, RW, and CAL at different time points.

RD decreased from 3.27 ± 0.70 mm at the baseline to 0.82 ± 0.60 mm at 2 months and 0.63 ± 0.40 mm at 6 months postoperatively. This reduction, according to Friedman test, was statistically significant ($P < 0.001$). The results of Wilcoxon signed-rank test showed that the reduction in RD between 2 and 6 months was not significant ($P = 0.083$), but the reduction in RD at 2 and 6 months compared to the baseline was statistically significant ($P = 0.001$).

At the baseline, RW was 2.67 ± 0.81 mm, which significantly decreased to 1.22 ± 0.93 mm at 2 months and to 1.20 ± 0.80 mm at 6 months postoperatively ($P < 0.001$). The change from 2 to 6 months was not significant ($P = 0.317$), but significant changes were noted at 2 months ($P = 0.001$) and at 6 months ($P < 0.001$) compared to the baseline. CAL also significantly decreased from 4.80 ± 1.20 mm at the baseline to 1.93 ± 1.22 mm at 2 months and 1.33 ± 0.97 mm at 6 months postoperatively ($P < 0.001$).

Table 1: Comparison of clinical parameters (mm) between the two groups at different time points

Clinical parameters	Case group	Control group	P-value*
Recession depth (RD)			
Baseline	3.27 ± 0.70	3.20 ± 0.77	0.855
2 months	0.60 ± 0.82	0.93 ± 1.22	0.262
6 months	0.40 ± 0.63	0.73 ± 0.96	0.163
Recession width (RW)			
Baseline	2.67 ± 0.81	2.47 ± 0.74	0.429
2 months	0.93 ± 1.22	1.00 ± 1.30	0.739
6 months	0.80 ± 1.20	1.00 ± 1.30	0.480
Clinical attachment level (CAL)			
Baseline	4.80 ± 1.20	4.67 ± 1.11	0.586
2 months	1.93 ± 1.22	2.40 ± 1.68	0.327
6 months	1.33 ± 0.97	1.80 ± 1.32	0.142
Probing depth (PD)			
Baseline	1.53 ± 0.64	1.47 ± 0.64	0.655
2 months	1.53 ± 0.51	1.60 ± 0.63	0.655
6 months	1.13 ± 0.35	1.27 ± 0.45	0.134

*: Wilcoxon signed-rank test

These improvements were significant at 2 and 6 months compared to the baseline ($P=0.001$) and also at the time interval between 2 and 6 months ($P=0.007$).

The same changes were recorded in the control group.

A significant reduction occurred in RD from 3.20 ± 0.77 mm at the baseline to 1.22 ± 0.93 mm at 2 months and 0.96 ± 0.73 mm at 6 months postoperatively ($P=0.001$). This reduction during the time interval between 2 and 6 months was not significant ($P=0.083$), but the reduction at 2 and 6 months compared to the baseline was statistically significant ($P=0.001$).

RW decreased from 2.47 ± 0.74 mm at the baseline to 1.30 ± 1.00 mm at 2 and 6 months ($P<0.001$); this reduction was statistically significant compared to the baseline ($P=0.001$).

A significant reduction occurred in CAL from 4.67 ± 1.11 mm at the baseline to 2.40 ± 1.68 mm at 2 months and 1.80 ± 1.32 mm at 6 months postoperatively ($P<0.001$). The reduction at 2 and 6 months was significant compared to the baseline ($P=0.001$), but the change during the time interval between 2 and 6 months was not statistically significant ($P=0.014$).

PD did not experience any significant change at the mentioned time points in any group ($P>0.05$). No significant difference was found in root coverage between the case and control groups postoperatively ($P>0.05$, Table 2).

Table 3 shows the frequency of complete root coverage achieved postoperatively in the two groups.

DISCUSSION

The current study showed that the use of Er,Cr:YSGG laser for root surface conditioning had no significant effect on the outcome of the treatment of gingival recession with SCTG. The mean CAL, RD, and RW in both groups significantly decreased during the study period. The mean root coverage in the case and control groups at 2 months postoperatively was 81% and 74%, respectively. These values were 87% and 80% at 6 months, respectively. Complete root coverage in both groups had a frequency of 60% at 2 months. This value in the case group increased to 67% at 6 months but remained unchanged in the control group. These findings are comparable to those of previous studies reporting a range of 64.7% to 97.3% for the mean root coverage and 18.1% to 96.1% for complete root coverage [16,21,29].

The commonly used modalities for the treatment of gingival recession during the 1960s and 1970s included free gingival graft [6] and pedicle grafts [8-10]. SCTG was used for this purpose in the early 1980s [7]. Several studies have reported the high success rate and high predictability of this approach [13-15]. Thus, SCTG was used for the treatment of gingival recession in this study.

The final goal of the treatment of gingival recession is to achieve complete root coverage to improve esthetics and eliminate tooth hypersensitivity [16,21,29]. Several conditioners have been used for this purpose, but controversy exists regarding their efficacy for the improvement of clinical parameters.

Table 2: Comparison of postoperative root coverage in the two groups

Clinical parameters	Case group	Control group	P-value*
Root coverage (mm)			
2 months	2.67 ± 1.11	2.27 ± 1.03	0.207
6 months	2.86 ± 0.99	2.46 ± 0.83	0.196
Root coverage (%)			
2 months	81.11 ± 27.18	74.44 ± 33.10	0.396
6 months	87.22 ± 20.62	80.00 ± 26.12	0.244

*: Wilcoxon signed-rank test

Table 3: Frequency of complete root coverage achieved postoperatively in the two groups

Complete root coverage	A N(%)	B N(%)	C N(%)
Case Group			
2 Months	9 (60)	3 (20)	3 (20)
6 Months	10 (66.6)	4 (26.7)	1 (6.7)
Control Group			
2 Months	9 (60)	2 (13.3)	4 (26.7)
6 Months	9 (60)	5 (33.3)	1 (6.7)

Level of root coverage (A=100% B=50-99% C=0-49%)

Some studies reported that conditioned root surfaces showed higher percentages of complete root coverage compared to unconditioned areas [30,31]. In contrast, the results of some other studies indicated no significant advantage for root conditioners [3,18,20]. Our results revealed that root surface conditioning by Er,Cr:YSGG laser did not improve the clinical results of SCTG.

Published clinical data regarding the results of the application of Er,Cr:YSGG laser for the treatment of gingival recession are not available. However, the application of laser has been recommended as an adjunct for this purpose. Clinical studies have demonstrated that laser is beneficial for improving the results of regenerative treatments since it reinforces the attachment of regenerated periodontal structures [32-34]. In an in-vitro study, Fekrazad et al [28] evaluated the efficacy of Er,Cr:YSGG laser for root surface conditioning compared to EDTA and no use of conditioner. They stated that laser irradiation had a higher potential for reinforcement of the attachment of fibroblasts, while EDTA caused no significant change in the results [28].

Er:YAG laser is a promising modality for periodontal treatment [35,36]. Clinical studies have reported the optimal efficacy of erbium laser for the treatment of periodontal pockets via surgical and non-surgical techniques [37,38] and also for root conditioning [39,40].

Changes caused by the thermomechanical effects

of Er:YAG laser on the root surface include a change in the microstructure as well as thermal alterations [34,41-43]. The changes occurred in surface microstructure are considered advantageous for the primary attachment of cells and tissues in the clinical setting and result in the better formation of fibrin and blood clots [41-43]. Due to high absorption in water, erbium lasers have a high power for ablation of dental hard tissues [34,35] without causing significant thermal complications such as carbonization, melting, or crack formation in the root structure, which are often seen following the use of CO₂ and Nd:YAG lasers [34,44]. Considering the biocompatibility of the surfaces lased with Er:YAG laser, several studies have shown the better attachment and faster proliferation of fibroblasts on these surfaces compared to mechanically debrided surfaces [17,45]. The Er,Cr:YSGG laser parameters used in this study included 2780 nm wavelength, 0.75 W power, H mode, and 20 Hz repetition rate, according to the recommendations of a previous study, to preserve the biocompatibility of lased root surfaces [28].

In contrast, some authors reported that root surfaces lased with Er:YAG laser showed significant micron-scale irregularities in vitro [41,43]. Fujii et al [41] demonstrated that lased root surfaces had a specific microstructure along with denatured collagen fibers. Regarding other lasers, Trylovich et al [46] reported that the application of Nd:YAG laser changed the

biocompatibility of root surfaces, making them unsuitable for the attachment of fibroblasts. Fayad et al [44] reported the complete absence of fibroblast attachment to root surfaces following the use of CO₂ laser.

Some recent studies indicated that the application of Er:YAG laser significantly improves a number of clinical parameters [17,37,38]. Dilsiz et al [3] discussed that the application of Er:YAG laser for root biomodification does not improve the results of SCTG. Bouchard et al [47] and Caffesse et al [18] reported that root surface conditioning with citric acid has no effect on the clinical results of SCTG. Our findings were in agreement with their results.

Further clinical studies are required to confirm the results of this study in larger study populations with longer follow-up periods.

CONCLUSION

According to the results of this study, root surface conditioning by Er,Cr:YSGG laser improved the mean root coverage and the percentage of complete root coverage. However, these changes were not statistically significant.

ACKNOWLEDGMENTS

We would like to extend our gratitude to the Deputy of Research at Hamadan University of Medical Sciences and the Dental Research Center for the financial support provided.

REFERENCES

- 1- The American Academy of Periodontology. Glossary of Periodontal Terms. 4th ed. Chicago, IL, USA: The American Academy of Periodontology; 2001:44.
- 2- Dilsiz A, Aydin T, Canakci V, Cicek Y. Root Surface Biomodification with Nd:YAG Laser for the Treatment of Gingival Recession with Subepithelial Connective Tissue Grafts. *Photomed Laser Surg.* 2010 Jun;28(3):337-43.
- 3- Dilsiz A, Aydin T, Yavuz MS. Root Surface Biomodification with an Er:YAG Laser for the

Treatment of Gingival Recession with Subepithelial Connective Tissue Grafts. *Photomed Laser Surg.* 2010 Aug;28(4):511-7.

- 4- Roman A, Soanca A, Kasaj A, Stratul SI. Subepithelial Connective Tissue Graft with or without Enamel Matrix Derivative for the Treatment of Miller Class I and II Gingival Recessions: A Controlled Randomized Clinical Trial. *J Periodontol Res.* 2013 Oct;48(5):563-72.
- 5- Kassab MM, Cohen RE. The Etiology and Prevalence of Gingival Recession. *J Am Dent Assoc.* 2003 Feb;134(2):220-5.
- 6- Miller PD Jr. Root Coverage Using a Free Soft Tissue Autograft Following Citric Acid Application. Part 1: Technique. *Int J Periodontics Restorative Dent.* 1982;2(1):65-70.
- 7- Langer B, Langer L. Subepithelial Connective Tissue Graft Technique for Root Coverage. *J Periodontol.* 1985 Dec;56(12):715-20.
- 8- Bernimoulin JP, Lüscher B, Mühlemann HR. Coronally Repositioned Periodontal Flap. *J Clin Periodontol.* 1975 Mar;2(1):1-13.
- 9- Grupe HE, Warren RF Jr. Repair of Gingival Defects by a Sliding Flap Operation. *J Periodontol.* 1956 Apr;27(2):92-5.
- 10- Cohen DW, Ross SE. The Double Papillae Repositioned Flap in Periodontal Therapy. *J Periodontol.* 1968 Mar;39(2):65-70.
- 11- Pini Prato G, Tinti C, Vincenzi G, Magnani C, Cortellini P, Clauser C. Guided Tissue Regeneration Versus Mucogingival Surgery in the Treatment of Human Buccal Gingival Recession. *J Periodontol.* 1992 Nov;63(11):919-28.
- 12- Tal H, Moses O, Zohar R, Meir H, Nemcovsky C. Root Coverage of Advanced Gingival Recession: A Comparative Study Between Acellular Dermal Matrix Allograft and Subepithelial Connective Tissue Grafts. *J Periodontol.* 2002 Dec;73(12):1405-11.
- 13- Griffin TJ, Cheung WS, Zavras AI, Damoulis PD. Postoperative Complications Following Gingival Augmentation Procedures. *J Periodontol.* 2006 Dec;77(12):2070-9.
- 14- Harris RJ. The Connective Tissue and Partial Thickness Double Pedicle Graft: A Predictable

- Method of Obtaining Root Coverage. *J Periodontol.* 1992 May;63(5):477-86.
- 15- Han JS, John V, Blanchard SB, Kowolik MJ, Eckert GJ. Changes in Gingival Dimensions Following Connective Tissue Grafts for Root Coverage: Comparison of Two Procedures. *J Periodontol.* 2008 Aug;79(8):1346-54.
- 16- Chambrone L, Chambrone D, Pustiglioni FE, Chambrone LA, Lima LA. Can Subepithelial Connective Tissue Grafts be Considered the Gold Standard Procedure in the Treatment of Miller Class I and II Recession-Type Defects? *J Dent.* 2008 Sep;36(9):659-71.
- 17- Schwarz F, Aoki A, Sculean A, Georg T, Scherbaum W, Becker J. In Vivo Effects of an Er:YAG Laser, an Ultrasonic System and Scaling and Root Planing on the Biocompatibility of Periodontally Diseased Root Surfaces in Cultures of Human PDL Fibroblasts. *Laser Surg Med.* 2003 Aug;33(2):140-7.
- 18- Caffesse RG, De LaRosa M, Garza M, Munne-Travers A, Mondragon JC, Weltman R. Citric Acid Demineralization and Subepithelial Connective Tissue Grafts. *J Periodontol.* 2000 Apr;71(4):568-72.
- 19- Kassab MM, Cohen RE, Andreana S, Dentino AR. The Effect of EDTA in Attachment Gain and Root Coverage. *Compend Contin Educ Dent.* 2006 Jun;27(6):353-60.
- 20- Bouchard P, Nilveus R, Etienne D. Clinical Evaluation of Tetracycline HCl Conditioning in the Treatment of Gingival Recessions. A Comparative Study. *J Periodontol.* 1997 Mar;68(3):262-9.
- 21- Oates TW, Robinson M, Gunsolley JC. Surgical Therapies for the Treatment of Gingival Recession. A Systematic Review. *Ann Periodontol.* 2003 Dec;8(1):303-20.
- 22- Sallum EA, Casati MZ, Caffesse RG, Funis LP, Nociti Júnior FH, Sallum AW. Coronally Positioned Flap with or without Enamel Matrix Protein Derivative for the Treatment of Gingival Recessions. *Am J Dent.* 2003 Oct;16(5):287-91.
- 23- Lacoste E, Martineau I, Gagnon G. Platelet Concentrates: Effects of Calcium and Thrombin on Endothelial Cell Proliferation and Growth Factor Release. *J Periodontol.* 2003 Oct;74(10):1498-507.
- 24- Rubins RP, Tolmie PN, Corsig KT, Kerr EN, Kim DM. Subepithelial Connective Tissue Graft with Growth Factor for the Treatment of Maxillary Gingival Recession Defects. *Int J Periodontics Restorative Dent.* 2013 Jan-Feb;33(1):43-50.
- 25- Kreisler M, Al Haj H, d'Hoedt B. Clinical Efficacy of Semiconductor Laser Application as an Adjunct to Conventional Scaling and Root Planing. *Laser Surg Med.* 2005 Dec;37(5):350-5.
- 26- Harashima T, Kinoshita J, Kimura Y, Brugnera A, Zanin F, Pecora JD, et al. Morphological Comparative Study on Ablation of Dental Hard Tissues at Cavity Preparation by Er:YAG and Er,Cr:YSGG Lasers. *Photomed Laser Surg.* 2005 Feb;23(1):52-5.
- 27- Miller PD Jr. A Classification of Marginal Tissue Recession. *Int J Periodontics Restorative Dent.* 1985;5(2):8-13.
- 28- Fekrazad R, Lotfi G, Harandi M, Ayremlou S, Kalhori KA. Evaluation of Fibroblast Attachment in Root Conditioning with Er,Cr:YSGG Laser Versus EDTA: A SEM Study. *Microsc Res Tech.* 2015 Apr;78(4):317-22.
- 29- Rocuzzo M, Bunino M, Needleman I, Sanz M. Periodontal Plastic Surgery for Treatment of Localized Gingival Recessions: A Systematic Review. *J Clin Periodontol.* 2002;29 Suppl 3:178-94; discussion 195-6.
- 30- Miller PD Jr. Root Coverage Using the Free Soft Tissue Autograft Following Citric Acid Application. III. A Successful and Predictable Procedure in Areas of Deep-Wide Recession. *Int J Periodontics Restorative Dent.* 1985;5(2):14-37.
- 31- Tolmie PN, Rubins RP, Buck GS, Vagianos V, Lanz JC. The Predictability of Root Coverage by way of Free Gingival Autografts and Citric Acid Application: An Evaluation by Multiple Clinicians. *Int J Periodontics Restorative Dent.* 1991;11(4):261-71.
- 32- Crespi R, Barone A, Covani U, Ciaglia RN, Romanos GE. Effects of CO2 Laser Treatment on Fibroblast Attachment to Root Surfaces: A Scanning Electron Microscopy Analysis. *J Periodontol.* 2002

- Nov;73(11):1308-12.
- 33- Pant V, Dixit J, Agrawal AK, Seth PK, Pant AB. Behavior of Human Periodontal Ligament Cells on CO₂ Laser Irradiated Dentinal Root Surfaces: An In Vitro Study. *J Periodontal Res.* 2004;39:373-9.
- 34- Israel M, Cobb CM, Rossmann JA, Spencer P. The Effects of CO₂, Nd:YAG and Er:YAG Lasers with and without Surface Coolant on Tooth Root Surfaces: An In Vitro Study. *J Clin Periodontol.* 1997 Sep;24(9 Pt 1):595-602.
- 35- Schwarz F, Aoki A, Becker J, Sculean A. Laser Application in Non-Surgical Periodontal Therapy: A Systematic Review. *J Clin Periodontol.* 2008 Sep;35(8 Suppl):29-44.
- 36- Ishikawa I, Aoki A, Takasaki AA. Potential Applications of Erbium:YAG Laser in Periodontics. *J Periodontal Res.* 2004 Aug;39(4):275-85.
- 37- Schwarz F, Sculean A, Berakdar M, Szathmari L, Georg T, Becker J. In Vivo and In Vitro Effects of an Er:YAG Laser, A GaAlAs Diode Laser, and Scaling and Root Planing on Periodontally Diseased Root Surfaces: A Comparative Histologic Study. *Lasers Surg Med.* 2003;32(5):359-66.
- 38- Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Becker J. Clinical Evaluation of an Er:YAG Laser Combined with Scaling and Root Planing for Non-Surgical Periodontal Treatment: A Controlled, Prospective Clinical Study. *J Clin Periodontol.* 2003 Jan;30(1):26-34.
- 39- Lavu V, Sundaram S, Sabarish R, Rao SR. Root Surface Bio-modification with Erbium Lasers- A Myth or a Reality?? *Open Dent J.* 2015 Jan 30;9:79-86.
- 40- Cekisi A, Maden I, Yildiz S, San T, Isik G. Evaluation of Blood Cell Attachment on Er:YAG Laser Applied Root Surface Using Scanning Electron Microscopy. *Int J Med Sci.* 2013;10(5):560-6.
- 41- Fujii T, Baehni PC, Kawai O, Kawakami T, Matsuda K, Kowashi Y. Scanning Electron Microscopic Study of the Effects of Er:YAG Laser on Root Cementum. *J Periodontol.* 1998 Nov;69(11):1283-90.
- 42- Theodoro LH, Sampaio JE, Haypek P, Bachmann L, Zezell DM, Garcia VG. Effect of Er:YAG and Diode Lasers on the Adhesion of Blood Components and on the Morphology of Irradiated Root Surfaces. *J Periodontal Res.* 2006 Oct;41(5):381-90.
- 43- Sasaki KM, Aoki A, Ichinose S, Ishikawa I. Morphological Analysis of Cementum and Root Dentin after Er:YAG Laser Irradiation. *Lasers Surg Med.* 2002;31(2):79-85.
- 44- Fayad MI, Hawkinson R, Daniel J, Hao J. The Effect of CO₂ Laser Irradiation on PDL Cell Attachment to Resected Root Surfaces. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004 Apr;97(4):518-23.
- 45- Mizutani K, Aoki A, Takasaki AA, Kinoshita A, Hayashi C, Oda S, et al. Periodontal Tissue Healing Following Flap Surgery Using an Er:YAG Laser in Dogs. *Lasers Surg Med.* 2006 Apr;38(4):314-24.
- 46- Trylovich DJ, Cobb CM, Pippin DJ, Spencer P, Killoy WJ. The Effects of the Nd:YAG Laser on In Vitro Fibroblast Attachment to Endotoxin-Treated Root Surfaces. *J Periodontol.* 1992 Jul;63(7):626-32.
- 47- Bouchard P, Etienne D, Ouhayoun JP, Nilvéus R. Subepithelial Connective Tissue Grafts in the Treatment of Gingival Recessions: A Comparative Study of 2 Procedures. *J Periodontol.* 1994 Oct;65(10):929-36.